

Nuclear modification of high- p_T hadron spectra in $p + A$ collisions at LHC

Rong Xu,¹ Wei-Tian Deng,² and Xin-Nian Wang^{3,4}

¹*Key Laboratory of Quark and Lepton Physics (MOE) and Institute of Particle Physics,
Central China Normal University, Wuhan 430079, China*

²*Frankfurt Institute for Advanced Studies, Ruth-Moufang-Strasse 1, D-60438 Frankfurt am Main, Germany*

³*Institute of Particle Physics, Central China Normal University, Wuhan 430079, China*

⁴*Nuclear Science Division, MS 70R0319, Lawrence Berkeley National Laboratory, Berkeley, California 94720*

Multiple parton scatterings in high-energy $p + A$ collisions involve multi-parton correlation inside the projectile and color coherence of multiple jets which will lead to nuclear modification of final hadron spectra relative to that in $p + p$ collisions. Such modifications of final hadron spectra in $p + A$ collisions are studied within the HIJING 2.1 model. Besides parton shadowing and transverse momentum broadening, which influence final hadron spectra at intermediate p_T , modification of parton flavor composition due to multiple scattering and hadronization of many-parton jets system are shown to lead to suppression of final hadrons at large p_T in $p + A$ collisions at the Large Hadron Collider.

PACS numbers: 24.85.+p, 25.75.-q, 25.75.Bh

In search for the formation of quark-gluon plasma (QGP) in high-energy heavy-ion collisions, jet quenching or suppression of high- p_T hadrons [1] can be used for tomographic study of medium properties [2, 3]. However, initial multiple parton scatterings in cold nuclei before the formation of QGP can also lead to nuclear modification of final high- p_T hadron spectra that needs to be understood for a quantitative study of QGP properties through jet quenching.

Parton shadowing or nuclear modification of parton distributions inside large nuclei is one of the most studied cold nuclear effects. The QCD evolution, however, dramatically reduces the effect on final high- p_T hadron spectra in $p + A$ collisions [4–7]. Modification of parton distributions in the projectile nucleon through parton energy loss [8, 9] is estimated to have a small effect on hadron spectra in the central rapidity region of high-energy $p + A$ collisions. These cold nuclear effects have also been studied within the Color Glass Condensate (CGC) model [10–12]. However, multiple parton correlations inside the projectile nucleon should also be considered. Energy-momentum and valence quark number conservation alone could modify the momentum and flavor dependence of final-state parton and hadron spectra at large p_T . Hadronization of multiple-jet systems from multiple scattering in $p + A$ collisions could also affect final hadron spectra at intermediate and large p_T .

In this paper, we study the nuclear modification of parton and hadron spectra due to parton shadowing, initial and final-state transverse momentum broadening, final-state parton flavor composition and hadronization of multiple jets within the HIJING [13] Monte Carlo model. It is based on a two-component model for hadron production in high-energy nucleon and nuclear collisions. The soft and hard components are separated by a cut-off p_0 in the transverse momentum exchange. Hard parton scatterings with $p_T > p_0$ are assumed to be described by the perturbative QCD (pQCD) and soft interactions are approximated by string excitations with an effective

cross section σ_{soft} . In HIJING 2.0 [14], parton distribution functions (PDF's) are given by the Gluck-Reya-Vogt (GRV) parameterization [15]. An energy-dependent cut-off $p_0(\sqrt{s})$ and soft cross section $\sigma_{\text{soft}}(\sqrt{s})$ are assumed in order to fit experimental values of the total and inelastic cross sections of $p + p(\bar{p})$ collisions and the energy-dependence of the central rapidity density of charged hadron multiplicities. HIJING 2.0 can describe most features of hadron production in $p + p$ collisions at colliding energies up to 7 TeV at the Large Hadron Collider (LHC) [14].

In $p + A$ and $A + A$ collisions, one has to consider nuclear modification of initial parton distributions due to multiple scatterings. HIJING 2.0 employs a factorized form of parton distributions in nuclei,

$$f_{a/A}(x, Q^2, b) = AR_a^A(x, Q^2, b)f_{a/N}(x, Q^2), \quad (1)$$

where Q is the momentum scale, b the impact-parameter, x the momentum fraction carried by the parton a and $R_a^A(x, Q^2, b)$ is the impact-parameter-dependent nuclear modification factor as given by the new HIJING parameterization [16]. It was fit to the centrality dependence of the pseudo-rapidity multiplicity density per participant pair $2dN_{ch}/d\eta/N_{\text{part}}$ at the Relativistic Heavy-ion Collider (RHIC), and also describes the data at the LHC energies as well [14].

In addition to modification of initial parton distributions, multiple scatterings inside a nucleus should also lead to transverse momentum (k_T) broadening of both initial and final-state partons. The k_T -broadening is responsible for the Cronin effect or enhancement of intermediate- p_T hadron spectra in $p + A$ collisions [17]. It is implemented in HIJING 2.1 as a k_T -kick by each binary nucleon-nucleon scattering to the initial partons of a hard scattering as well as final-state partons produced prior to the current scattering. The k_T -kick for each scattering follows a Gaussian distribution and an energy dependence of the width,

$$\langle k_T^2 \rangle = 0.14 \log(\sqrt{s}/\text{GeV}) - 0.43 \text{ GeV}^2, \quad (2)$$




FIG. 1: (color online) Charged hadron rapidity distribution from HIJING 2.1 with (solid) and without (dot-dashed) parton shadowing in $p(d) + A$ and $p + p$ (dashed) collisions at $\sqrt{s_{NN}} = 200$ GeV (upper) and 5.0 TeV (lower) as compared to data from the STAR [18] and PHOBOS [19] experiment.

is assumed to describe the measured Cronin effect in $p + A$ collisions. After the k_T -kick, longitudinal momenta of partons are reshuffled pair-wise between projectile and target partons in order to ensure four-momentum conservation. Consequently, introduction of the k_T -kick will also influence hadron rapidity distributions. We have to fine-tune the gluon shadowing parameter s_g [14, 16] again to describe experimental data on rapidity distributions.

Shown in Fig. 1 are HIJING 2.1 results on rapidity distributions of charged hadrons in minimum-biased $d + Au$ (upper panel) and $p + Pb$ (lower panel) collisions (solid) at $\sqrt{s_{NN}} = 200$ GeV and 5.0 TeV, respectively, as compared to $d + Au$ data from the STAR experiment [18]. Here the gluon shadowing parameter is set to $s_g = 0.28$. Also shown are HIJING2.1 results for $p + p$ collisions (dashed) as compared to the PHOBOS data [19] and $p(d) + A$ collisions without parton shadowing (dot-dashed). Gluon shadowing in HIJING 2.1 is seen to suppress the hadron yield in mid and forward rapidity region, especially at the LHC energy. Measurement of the hadron rapidity distribution at LHC therefore would provide further constraints on the gluon shadowing.

The hard processes in HIJING are based on lowest order pQCD in which the single jet inclusive cross section,

$$\frac{d\sigma_{pA}^{jet}}{dy_1 d^2p_T} = K \int dy_2 d^2b t_A(b) \sum_{a,b,c} x_1 f_{a/p}(x_1, p_T^2) \times x_2 f_{a/A}(x_2, p_T^2, b) \frac{d\sigma_{ab \rightarrow cb}}{dt}, \quad (3)$$

is directly proportional to nuclear parton distributions




FIG. 2: (color online) Nuclear modification factor for parton p_T spectra in $p(d) + A$ collisions at $\sqrt{s_{NN}} = 200$ GeV (upper) and 5.0 TeV (lower) from HIJING 2.1 with (solid), without (dashed) parton shadowing and with de-coherent hard scatterings (DHC) (dot-dashed and dotted).

$f_{a/A}(x_2, Q^2, b)$. Here $x_{1,2} = p_T(e^{\pm y_1} + e^{\pm y_2})/\sqrt{s}$ are fractional momenta of the initial partons, $y_{1,2}$ are rapidities of the final-state parton jets and $K \approx 2$ accounts for higher-order corrections. The nuclear thickness function is normalized to $\int d^2b t_A(b) = 1$. In the central rapidity region ($y_{1,2} = 0$), transverse momenta of final-state jets determine the momentum fractions $x_{1,2} = 2p_T/\sqrt{s}$ of the initial colliding partons. Therefore, p_T spectra of produced partons via hard scatterings in $p + A$ collisions should depend on the nuclear modification of parton distributions in nuclei. To study parton shadowing and other cold nuclear effects, we calculate the nuclear modification factor for final-state parton and hadron p_T spectra,

$$R_{pA}(p_T) = \frac{dN_{pA}/dy d^2p_T}{\langle N_{bin} \rangle dN_{pp}/dy d^2p_T}, \quad (4)$$

where $\langle N_{bin} \rangle$ is the average number of binary nucleon-nucleon interactions in $p + A$ collisions.

Shown in Fig. 2 are parton modification factors from the HIJING 2.1 for minimum-biased $d + Au$ and $p + Pb$ collisions at $\sqrt{s_{NN}} = 200$ GeV and 5.0 TeV, respectively. The enhancement of parton spectra at intermediate p_T is the signature effect of k_T -broadening, which should disappear at large p_T . In the HIJING model, hard scatterings are simulated each time the projectile parton interact with a target parton alongside soft interactions with the remnants of colliding nucleons. HIJING ensures energy-momentum conservation by subtracting




FIG. 3: (color online) Quark to gluon ratio as a function of p_T in $p + p$ (dotted) $p(d) + A$ collisions at the RHIC (upper) and LHC (lower) energies from HIJING 2.1 without (solid) and with (dashed) de-coherent hard scatterings (DHC).

energy-momentum transfers in the previous hard and soft interactions from the projectile and thus restricting energy available for subsequent hard collisions. Since the time scale of hard scatterings is much shorter than soft interactions, such a coupling between hard and soft interactions might not be physical. As an option in our study, we turn off such a coupling and do not restrict energy available for hard scatterings. We denote the results as DHC (de-coherent hard scattering). One can see from Fig. 2, both parton shadowing and soft-hard coupling suppresses high- p_T parton production. These features of parton spectra will be reflected in the final hadron spectra after hadronization.

In $p + A$ collisions, the projectile proton scatters with multiple nucleons inside the target, each with finite probability of hard parton scattering. The leading contribution comes from independent hard scatterings, each involving independent initial partons from projectile and target nucleons. In principle, a fast projectile parton can also suffer multiple hard scatterings which will affect the final-state parton spectra but do not increase the projectile's contribution to the final-state parton multiplicity. Momentum correlation of multiple partons inside the projectile is neglected beyond the conservation of the total momentum. Flavor conservation, however, will limit the availability of valence quarks from the projectile for each of these multiple independent hard scatterings as the projectile plunges through the target nucleus. Such limitation on the number of valence quarks available for




FIG. 4: (color online) Proton to pion ratio at RHIC and LHC from HIJING 2.1 with different options: default (solid), without shadowing (ashed), DHC (dot-dashed), DHC without parton shadowing (dotted) in $d + Au$ collisions at RHIC and $p + Pb$ at LHC. The ratio in $p + p$ collisions is also shown (long dashed). The RHIC experimental data are from STAR [21].

each independent hard scattering will lead to a change in the flavor composition of produced partons per average binary nucleon-nucleon collision. Shown in Fig. 3 are the ratios of produced quark and gluon p_T spectra in the central rapidity region of $d + Au$ and $p + Pb$ collisions at $\sqrt{s}_{NN} = 200$ GeV and 5.0 TeV, respectively, as compared to that in $p + p$ collisions. The quark number conservation in multiple scatterings is seen to lead to smaller quark/gluon ratio in $p + A$ relative to $p + p$ collisions. The suppressed quark yield and high quark/gluon ratio, especially at RHIC, will lead to the modification of final-state parton spectra at large p_T as shown in Fig. 2. Because gluon fragmentation functions are softer than quarks, the increased fraction of gluon jets in $p + A$ collisions will lead to apparent suppression of high p_T hadron spectra relative to $p + p$ collisions.

The modified flavor composition of the final parton due to multiple scattering in pA collisions could also change the flavor composition of final hadrons due to flavor-dependence of the fragmentation functions. Gluon jets in general have higher proton-to-pion ratio than in quark jets. HIJING has a strong gluon shadowing which suppresses larger p_T gluon jets. This should reduce the proton-to-pion ratio in HIJING 2.1 results as compared to the results without gluon shadowing as shown in Fig. 4. This reduction of proton-to-pion ratio by gluon shadow-

ing is much stronger at the LHC energy. Nevertheless, the proton-to-pion ratio in $d + Au$ collisions at RHIC and $p + Pb$ collisions at LHC will still be larger than that in $p + p$ collisions because of the enhanced gluon-to-quark ratio in $p + A$ collisions relative to that in $p + p$ as shown in Fig. 3. Such a trend is already seen in the STAR data [21] at RHIC. The flavor composition of final hadrons in $p + A$ collisions could also be influenced by other non-perturbative mechanisms such as parton recombination during hadronization [22] and hadronic final-state rescattering [23].

Another process that can also modify final hadron spectra in $p + A$ collisions is jet hadronization or fragmentation. HIJING follows the same routine as in PYTHIA that groups jet partons together with project and target remnants to form color-singlet string systems. In HIJING, jet parton showers from all hard scatterings, including initial and final-state radiations, are ordered in rapidity and gluons are then connected to the valence quark and diquark of the associated projectile or target nucleons as kinks to form string systems. These string systems are hadronized according to the Lund string model [20]. The projectile can undergo multiple scatterings in $p + A$ collisions. Its string system therefore has more gluons attached as compared to that in $p + p$ collisions. Hadrons from such a string system are expected to be softer than independent fragmentation of individual gluons. To test the sensitivity of nuclear modification of final hadron spectra to jet hadronization, we also consider the independent fragmentation mechanism in HIJING for both $p + p$ and $p + A$ collisions.

Shown in Fig. 5 are nuclear modification factors for charged hadrons in $d + Au$ and $p + Pb$ collisions at $\sqrt{s_{NN}} = 200$ GeV and 5.0 TeV, respectively, as compared to existing data from both the PHENIX and STAR expok, eriment at RHIC [24, 25]. At low p_T , hadron production is dominated by hadronization of soft strings which is proportional to the number of participant nucleons $N_{part} = 1 + N_{bin}$. The nuclear modification factor should approach $R_{pA} \approx (1 + 1/N_{bin})/2 \approx 1/2$ in the limit $N_{bin} \gg 1$. At large p_T , incoherent hard parton scatterings become dominate and $R_{pA} \approx 1$ without any nuclear modification. The peak in the nuclear modification factor is generally known as the Cronin enhancement of hadron spectra at intermediate p_T due to k_T -broadening through multiple scatterings. Such a peak has been observed in pA collisions at the Fermilab energies ($E_{lab} = 200 - 800$ GeV) [26] and the position of the peak at around 4 GeV/c has no apparent energy dependence and remains the same in HIJING results at both RHIC and LHC energies. Nuclear shadowing of parton distributions at LHC will only affect the overall value of R_{pA} at intermediate p_T , but the peak-feature and its position remain the same. This is a unique prediction of the k_T -broadening due to multiple scattering as implemented in HIJING2.1. It is in sharp contrast to the prediction of disappearance of the peak in gluon saturation model or relocation to much higher p_T due to strong




FIG. 5: (color online) Nuclear modification factor for charged hadrons in $p + A$ collisions from HIJING 2.1 with different options: default (solid), DHC (dashed), DHC without parton shadowing (dot-dashed) and DHC without parton shadowing and independent fragmentation (dotted), as compared to data from the PHENIX and STAR experiment [24, 25]. The arrowed lines indicate the most possible behavior of the nuclear modification factors at LHC between low and high p_T regions.

anti-shadowing in the collinear pQCD parton model. Experimental test of these features will shed light on the parton dynamics in high-energy nuclei.

At very large p_T , parton shadowing in nuclei, soft-hard coupling and enhanced gluonic jets due to valence quark conservation all lead to suppression of charged hadron spectra. Fragmentation of the projectile string system with multiple gluon jets in $p + A$ collisions seems to further suppress high- p_T hadron spectra as compared to the independent fragmentation (dotted). These nuclear effects are consistent with existing $d + Au$ data at RHIC. However, $p + Pb$ collisions at LHC will be able to test these scenarios within the available p_T range. Since parton shadowing will disappear at large p_T [4] due to QCD evolution and hard scatterings are de-coherent from soft interactions, the nuclear modification factor at LHC will likely follow the default and DHC results at low p_T and approach DHC without shadowing at large p_T with possible further modifications due to hadronization of multiple jets. In Fig. 5, we indicate this trend by joining these two set of results with arrowed lines at intermediate p_T .

The initial-state interaction and fragmentation of multiple jet systems should also affect the final hadron spectra in high-energy heavy-ion collisions in addition to jet quenching due to final-state interaction. The existing ex-




FIG. 6: (color online) Nuclear modification factors for final parton (upper panel) and charged hadrons (lower panel) in central Pb+Pb collisions at $\sqrt{s} = 2.76$ TeV from HIJING 2.1 with different options: default (solid), DHC (dashed), DHC without parton shadowing (dot-dashed) and DHC without parton shadowing and independent fragmentation (dotted), as compared to data from the ALICE and CMS experiment [27, 28]. Jet quenching due to final-state interaction is not considered in the HIJING calculations.

perimental data on hadron spectra modification should provide some constraints on the initial-state parton interactions and the final hadronization mechanism. Shown in Fig. 6 are nuclear modification factors for final parton (upper panel) and hadron spectra in central $Pb+Pb$ collisions at $\sqrt{s} = 2.76$ TeV. Note that there is no jet quenching effect in the calculation shown. On the parton level, the nuclear modification at large p_T mainly comes from parton shadowing which should be small if the scale dependence of shadowing is taken into account. The final hadron spectra at large p_T are however significantly suppressed due to the Lund string fragmentation of multiple parton systems (solid and dashed curves in the lower panel) leaving no room for additional suppression by jet quenching as compared to experimental data from ALICE [27] and CMS [28]. This is a strong indication that the Lund string fragmentation for hadroniza-

tion is no longer a valid mode for multiple gluonic jet systems in heavy-ion collisions where a deconfined quark-gluon plasma is formed. Recent studies show that multiple parton scatterings inside a quark-gluon plasma can in fact change the color flow of the jet parton shower from that in vacuum [29–32]. One therefore should expect the fragmentation processes in heavy-ion collisions to be significantly modified from that in vacuum. An independent jet fragmentation mechanism such as parton recombination model might be a better alternative where local parton phase distributions determine the final hadron spectra and global color flow of the multiple jet system is no longer important. Since multiple gluonic jet systems in pA collisions can be considered as precursors of the quark-gluon plasma, investigation of the final hadron spectra at large p_T in $p+A$ collisions can shed light on the hadronization mechanism of energetic jets after they escape jet quenching and emerge from the quark-gluon plasma in high-energy heavy-ion collisions.

In summary, we have studied the nuclear modification of hadron spectra in $p+A$ collisions at both the RHIC and LHC energies within HIJING2.1 Monte Carlo model and their sensitivities to cold nuclear effects. Multiple parton correlations in the projectile due to valence quark number conservation, string fragmentation of multiple gluon jets as well as parton shadowing can all lead to suppression of high- p_T hadron spectra in $p+A$ relative to $p+p$ collisions. Comparisons between HIJING2.1 results without jet quenching and existing data on large p_T hadron suppression at LHC indicate an independence fragmentation mechanism for multiple jet systems such as in pA and AA collisions. Experimental test of these effects in $p+Pb$ collisions at LHC and its implications for hard scatterings in $A+A$ collisions is crucial for the study of jet quenching and extraction of jet transport properties in the hot quark-gluon plasma.

Acknowledgement

We thank M. Gyulassy for helpful discussions. This work was supported in part by the NSFC under the project No. 10825523 and 11221504, by self-determined research funds of CCNU from the colleges' basic research and operation of MOE, Helmholtz International Center for FAIR within the framework of the LOEWE program launched by the State of Hesse, and US Department of Energy under Contract No. DE-AC02-05CH11231 and within the framework of the JET Collaboration.

-
- [1] X. N. Wang and M. Gyulassy, Phys. Rev. Lett. **68**, 1480 (1992).
 - [2] M. Gyulassy, P. Levai and I. Vitev, Phys. Lett. B **538**, 282 (2002).
 - [3] E. Wang and X. -N. Wang, Phys. Rev. Lett. **89**, 162301

- (2002).
- [4] K. J. Eskola, H. Paukkunen and C. A. Salgado, JHEP **0904**, 065 (2009)
- [5] I. Vitev, M. Gyulassy, Phys. Rev. Lett. **89**, 252301 (2002).

- [6] P. Quiroga-Arias, J. G. Milhano, U. A. Wiedemann, Nucl. Phys. **A855**, 498-501 (2011).
- [7] G. G. Barnafoldi, J. Barrette, M. Gyulassy, P. Levai, V. Topor Pop, arXiv:1111.3646.
- [8] R. B. Neufeld, I. Vitev, B. -W. Zhang, Phys. Lett. **B704**, 590-595 (2011).
- [9] H. Xing, Y. Guo, E. Wang and X. -N. Wang, Nucl. Phys. A **879**, 77 (2012).
- [10] L. D. McLerran and R. Venugopalan, Phys. Rev. D **49**, 2233 (1994).
- [11] D. Kharzeev, Y. V. Kovchegov and K. Tuchin, Phys. Rev. D **68**, 094013 (2003).
- [12] J. Jalilian-Marian and A. H. Rezaeian, Phys. Rev. D **85**, 014017 (2012).
- [13] X. N. Wang and M. Gyulassy, Phys. Rev. D **44**, 3501 (1991); M. Gyulassy and X. N. Wang, Comput. Phys. Commun. **83**, 307 (1994).
- [14] W. -T. Deng, X. -N. Wang, R. Xu, Phys. Rev. **C83**, 014915 (2011); Phys. Lett. **B701**, 133-136 (2011).
- [15] M. Gluck, E. Reya and A. Vogt, Z. Phys. C **67**, 433 (1995).
- [16] S. -Y. Li, X. -N. Wang, Phys. Lett. **B527**, 85-91 (2002).
- [17] J. W. Cronin *et al.*, Phys. Rev. D **11**, 3105 (1975).
- [18] B. I. Abelev *et al.*, arXiv:nucl-ex/0703016.
- [19] B. Alver *et al.*, Phys. Rev. C **83**, 024913 (2011).
- [20] B. Andersson, G. Gustafson, G. Ingelman, T. Sjostrand, Phys. Rept. **97**, 31-145 (1983).
- [21] J. Adams *et al.* [STAR Collaboration], Phys. Lett. B **637**, 161 (2006) [nucl-ex/0601033].
- [22] R. C. Hwa and C. B. Yang, Phys. Rev. Lett. **93**, 082302 (2004).
- [23] B. Z. Kopeliovich, Phys. Lett. **B243**, 141 (1990).
- [24] S. S. Adler *et al.*, Phys. Rev. Lett. **91**, 072303 (2003).
- [25] J. Adams *et al.*, Phys. Rev. Lett. **91**, 072304 (2003).
- [26] See Fig. 9 in Phys. Rev. C **61**, 064910 (2000) and reference therein.
- [27] K. Aamodt *et al.* [ALICE Collaboration], Phys. Lett. B **696**, 30 (2011) [arXiv:1012.1004 [nucl-ex]].
- [28] S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **72**, 1945 (2012).
- [29] A. Leonidov and V. Nechitailo, Eur. Phys. J. C **71**, 1537 (2011) [arXiv:1006.0366 [nucl-th]].
- [30] J. Casalderrey-Solana and E. Iancu, JHEP **1108**, 015 (2011) [arXiv:1105.1760 [hep-ph]].
- [31] Y. Mehtar-Tani, C. A. Salgado and K. Tywoniuk, Phys. Rev. Lett. **106**, 122002 (2011) [arXiv:1009.2965 [hep-ph]].
- [32] A. Beraudo, J. G. Milhano and U. A. Wiedemann, JHEP **1207** (2012) 144 [arXiv:1204.4342 [hep-ph]].